arXiv:1211.3241v1 [quant-ph] 14 Nov 2012

# Genuine Multipartite Entanglement Trends in Gapless-gapped Transitions of Quantum Spin Systems

Anindya Biswas, R. Prabhu, Aditi Sen(De), and Ujjwal Sen
*Harish-Chandra Research Institute, Chhatnag Road, Jhunsi, Allahabad 211 019, India*

We investigate the behavior of a genuine multiparticle entanglement, viz. the generalized geometric measure, in gapless-to-gapped quantum transitions of one-dimensional and two-dimensional spin models. The investigations are performed in the exactly solvable one-dimensional $XY$ models, as well as two-dimensional frustrated $J_1 - J_2$ models, including the Shastry-Sutherland model. In particular, we find that multiparty entanglement acts as a better tool than the bipartite one in detecting such transitions in the frustrated quantum systems.

## I. INTRODUCTION

Recent developments at the interface of quantum information science and many-body physics indicate that quantum correlations, in particular, quantum entanglement [1] can potentially be a universal physical characteristic to investigate many-body phenomena. For example, entanglement has been proposed as a detector of quantum phase transitions in spin systems [2–4] and as a tool to develop and simplify efficient numerical simulations like density matrix renormalization group [5]. Of late, the developments involving the numerical simulations using matrix product states [6], projected entangled pair states [7] and tensor network states [8], have established a strong connection between entanglement and many-body theory. At the same time, the behavior of entanglement has been investigated in many-body systems like cold atomic gases in optical lattices [9], and trapped gaseous Bose-Einstein condensates [10]. Characterization of entanglement in such systems is mostly restricted to bipartite entanglement due to the general unavailability of computatable measures in the multipartite scenario. However, in some cases, it turns out that the bipartite entanglement measures can not capture the co-operative phenomena in the system [11, 12], and therefore, it is natural to look out for multipartite entanglement measures to investigate such many-body systems [13].

Multipartite entanglement measures, e.g. the geometric measure [14] (cf. [15]) and global entanglement [16] have been used to describe many-body phenomena [17]. However, in general, they are hard to compute and it is therefore not possible to use them for states of arbitrary many-body systems. Recently, a genuine multiparticle entanglement measure called generalized geometric measure (GGM) [18] has been introduced which can be easily computed for pure states in arbitrary dimensions and of an arbitrary number of particles. It has since been found to be useful to study the genuine multiparty entanglement present in systems like resonating valence bond states and states of disordered systems [19]. In this paper, we apply the GGM to study quantum spin models, including frustrated ones [20, 21]. Frustration appears in a many-body system if it is not possible to simultaneously and independently minimize all the interaction terms of the corresponding Hamiltonian [21]. The characterization of such systems is typically hard to achieve [21]. We consider the following four classes of quantum spin systems –

1. the quantum one-dimensional (1D) $XY$ model [22],
2. the quasi-2D antiferromagnetic $J_1 - J_2$ Heisenberg model [23–25],
3. the antiferromagnetic $J_1 - J_2$ model on a 2D lattice [20], and
4. the Shastry-Sutherland model [26].

The choice of above models is due to their immense importance in understanding the different exotic phases in many-body systems including high-$T_c$ superconductivity [27]. Moreover, the recent experimental realizations of such spin models in the laboratories [3, 28–30], for example, in optical lattice [3, 31], trapped ions [32], photons [33], and nuclear magnetic resonance [34], have led to the interesting possibility of the observation of the many-body effects described here in the laboratories. Towards the unfolding of such many-body effects, we apply the multiparty entanglement measure, the GGM, to detect the transition from gapless to gapped phases and vice-versa in these models.

The paper is organized as follows: In Sec. II, we present a formal definition of the genuine multiparty entanglement measure, viz., the GGM, and we express the same in terms of easily computable Schmidt coefficients. The results are presented in Sec. III, where in each subsection, we begin with a brief description of one of the quantum spin models. Sections III A and III B deal, respectively, with the anisotropic XY and the $J_1 - J_2$ models, both in one dimension. In Sections III C and III D, we study two dimensional models – the 2D $J_1 - J_2$ model and the Shastry-Sutherland model. Finally, we draw our conclusions in Sec. IV.

## II. GENERALIZED GEOMETRIC MEASURE

In this section, we present a brief description of the GGM, and show that it is efficiently computable for pure quantum states of an arbitrary number of parties. We now define the genuine multipartite entanglement measure, viz. the GGM. A pure quantum state $|\psi\rangle_{A_1A_2\ldots A_N}$, shared between $N$ parties, $A_1, A_2, \ldots, A_N$, is said to be genuinely $N$-party entangled, if it is not a product across any bipartite partition. The GGM of a pure quantum state $|\psi\rangle_{A_1A_2\ldots A_N}$ is defined as

$$\mathcal{E}(|\psi\rangle_{A_1A_2\ldots A_N}) = 1 - \Lambda^2_{\max}(|\psi\rangle_{A_1A_2\ldots A_N}). \qquad (1)$$

Here $\Lambda_{\max}(|\psi\rangle_{A_1A_2\ldots A_N}) = \max|\langle\phi|\psi\rangle_{A_1A_2\ldots A_N}|$ where the maximization is taken over all pure states $|\phi\rangle$ which are not genuinely $N$-party entangled.

Let us enumerate some of the properties of GGM.

1. $\mathcal{E}$ is non-vanishing for all genuine multiparty entangled states, and vanishing for others.
2. $\mathcal{E}$ is monotonically non-increasing under local (quantum) operations and classical communication. (The proof follows from the theorem stated below and Ref. [35].)

We now state a proposition, which shows how the GGM can be expressed in terms of Schmidt coefficients.

**Theorem:** The generalized geometric measure (GGM) can be expressed as

$$\mathcal{E}(|\psi\rangle) = 1 - \max\{\lambda^2_{\mathcal{A}:\mathcal{B}} | \mathcal{A}\cup\mathcal{B} = \{A_1, \ldots, A_N\}, \mathcal{A}\cap\mathcal{B} = \emptyset\}, \qquad (2)$$

where $\lambda_{\mathcal{A}:\mathcal{B}}$ is the maximal Schmidt coefficient of $|\psi\rangle_{A_1A_2\ldots A_N}$ in the $\mathcal{A} : \mathcal{B}$ bipartite split.

## III. GGM DETECTS GAPLESS - GAPPED QUANTUM TRANSITIONS

In this section, we consider a series of paradigmatic quantum spin systems. They are taken up one by one in the different subsections. Each subsection begins with a brief description of the Hamiltonian corresponding to the quantum system under study. Subsequently, we study the behavior of the ground state of these models, and investigate the advantage of considering the genuine multipartite entanglement measure in these models.

### A. Anisotropic XY model

The Hamiltonian for the one-dimensional anisotropic quantum $XY$ model of $N$ quantum spin-half particles, arranged in an 1D array, is given by [22]

$$H_{XY} = \frac{J}{2}\left(\sum_{i=1}^{N}(1+\gamma)\sigma_i^x\sigma_{i+1}^x + (1-\gamma)\sigma_i^y\sigma_{i+1}^y\right) + h\sum_{i=1}^{N}\sigma_i^z, \qquad (3)$$

where $J$ is the coupling constant for the nearest neighbor interaction, $\gamma \in (0,1]$ is the anisotropy parameter, $\sigma$'s are the Pauli spin matrices, and $h$ represents the external transverse magnetic field applied across the system. In all the models considered in this paper, we impose the periodic boundary condition. The quantum XY Hamiltonian can be diagonalized by applying Jordan-Wigner, Fourier, and Bogoliubov transformations [22]. At zero temperature, it undergoes a quantum phase transition driven by the transverse magnetic field at $\lambda \equiv h/J = 1$. Moreover, it is also known that the model is gapped for all field strengths except at the point where the quantum phase transition occurs. Such transitions have been detected by using bipartite entanglement measures like concurrence and multipartite entanglement measure like geometric measure [17, 36]. We investigate the behavior of the genuine multipartite entanglement measure viz., the GGM, of the ground state, when it crosses from one gapped phase to another, through the gapless point.

The ground state of the system represented by the quantum XY Hamiltonian, as given in Eq. (3), can be analytically obtained by using Majorana fermions and it is also possible to get the eigenvalues of the local states corresponding to the ground state, in different bipartitions [4, 22] which are given by

$$e_{x_1x_2\ldots x_l} = \prod_{i=1}^{L}\frac{1+(-1)^{x_i}\nu_i}{2}, \quad x_i = 0, 1\ \forall i, \qquad (4)$$

where $\nu_i$'s are the eigenvalues of $G_L$. The latter is in turn given by

$$G_L = \begin{bmatrix} g_0 & \cdot\ \cdot & g_{L-1} \\ \cdot & \cdot\ \cdot & \cdot \\ -g_{L-1} & \cdot\ \cdot & g_0 \end{bmatrix}, \qquad (5)$$

with

$$g_l = \frac{1}{2\pi}\int_0^{2\pi} d\phi\, e^{-il\phi}\frac{\cos\phi - \lambda - i\gamma\sin\phi}{|\cos\phi - \lambda - i\gamma\sin\phi|}. \qquad (6)$$

In Fig. 1, we have plotted the GGM and the derivative of the GGM for the ground state of the $XY$ model with respect to the transverse magnetic field, $\lambda$, for different values of the anisotropic constant $\gamma$. The logarithmic divergence of the derivative of GGM captures the presence of the quantum phase transition at $\lambda = 1$. When $\gamma = 1$, which corresponds to the Ising model, the genuine multipartite entanglement is maximum when compared to the systems with lower values of $\gamma$.

### B. Quasi-2D Frustrated $J_1 - J_2$ Model

We will now consider the frustrated quasi two-dimensional $J_1 - J_2$ Heisenberg model, in which the nearest neighbor couplings, $J_1$, and the next-nearest neighbor couplings, $J_2$, are both antiferromagnetic. It was found that solid state systems like $SrCuO_2$ can be described by this model [37]. Moreover, advances in the field of cold atomic systems promise to create and control such models in the laboratory [3]. The Hamiltonian of this model,



FIG. 1. (Color online.) GGM of the transverse XY model. The GGM (actually $\mathcal{E} - 1/2$) and its derivative (both dimensionless) are plotted on the vertical axis for the anisotropic transverse XY model for different anisotropy parameters $\gamma$, against the dimensionless system parameter $\lambda$ on the horizontal axis. The dashed (blue), circled (pink), and dotted (green) lines are respectively for the Ising ($\gamma = 1$), $\gamma = 0.8$, and $\gamma = 0.2$ models. The derivatives of GGM diverges at the quantum critical point $\lambda = 1$.

with $N$ lattice sites on a chain, can be written as

$$H_{1D} = J_1 \sum_{i=1}^{N} \vec{\sigma}_i \cdot \vec{\sigma}_{i+1} + J_2 \sum_{i=1}^{N} \vec{\sigma}_i \cdot \vec{\sigma}_{i+2}, \qquad (7)$$

where $J_1$ and $J_2$ are both positive. In the parameter space, $\alpha \equiv J_2/J_1 = 0.5$ is known as the Majumdar-Ghosh point, and the system is highly frustrated there. For an even number of sites, the ground state at the Majumdar-Ghosh point is doubly degenerate, and the ground state manifold is spanned by the two dimers $|\psi^{\pm}_{MG}\rangle = \Pi_{i=1}^{N/2}(|0\rangle_{2i}|1\rangle_{2i\pm1} - |1\rangle_{2i}|0\rangle_{2i\pm1})$. Note that the model is gapped at this point [23]. For $\alpha = 0$, the Hamiltonian reduces to the spin$-\frac{1}{2}$ Heisenberg antiferromagnet and hence the ground state, which is a spin fluid state having gapless excitations [38], can be studied by Bethe ansatz. At other points, the ground state and the energy gap of this model were considered by using exact diagonalization, density matrix renormalization group method, bosonization technique, etc [25]. It is known that at $\alpha \approx 0.2411$, a phase transition from fluid to dimerization occurs [39]. In the weakly frustrated regime, $0 < \alpha \lesssim 0.24$, the system is gapless, and therefore critical [23, 24]. The system enters a dimerized regime, and is gapped, for higher values of the coupling parameter. It is known that bipartite entanglement cannot detect this gapless to gapped transition [12].

The multisite entanglement measure, GGM, is however able to detect the same. We perform exact diagonalization of the Hamiltonian using TITPACK ver. 2 developed by H. Nishimori [40] to find the ground state and then compute the GGM. In Fig. 2(a), a discontinuity in $\mathcal{E}$ at $\alpha = 0.5$ is visible, which signals the Majumdar-Ghosh point. It is also clear from Fig. 2(a) that the GGM changes its curvature at the gapless to gapped transition. To identify the exact phase transition point, we introduce an order parameter by using GGM ($\mathcal{E}$), which we call the bound GGM, defined as

$$\mathcal{E}_B = \mathcal{E} - \frac{d^2\mathcal{E}}{d\alpha^2}. \qquad (8)$$

We find that the fluid-dimer transition can be detected by the vanishing of the bound GGM at $\alpha \approx 0.23$. The discontinuities in the GGM curves, apart from the discontinuity at $\alpha = 0.5$, are arguably due to the finite and small system sizes. Note that the GGM curves asymptotically go to zero since the spin chain decouples into two spin chains with nearest neighbor interaction couplings, $J_2$, for very high values of the driving parameter $\alpha$. It may be noted here that although $\mathcal{E}_B = 0$ is a simple differential equation, its solution can be found only by knowing the relevant boundary conditions, which, in this case, can be thought of as being provided by the Hamiltonian corresponding to the system. This renders the differential equation intractable (analytically or numerically), and we do not follow that path.



(a)

(b)

FIG. 2. (Color online.) GGM and bound GGM for the quasi-2D frustrated antiferromagnet. In (a), the GGM (dimensionless) is plotted on the vertical axis and system parameter $\alpha$ (dimensionless) is plotted along horizontal axis. The change in curvature between $\alpha = 0.2$ and $\alpha = 0.4$ indicates the gapless to gapped phase transition. In (b), the bound GGM (dimensionless) is plotted on the vertical axis and system parameter $\alpha$ (dimensionless) is plotted along horizontal axis. The fluid-dimer transition is signaled by the vanishing of the bound GGM.



FIG. 3. (Color online.) GGM for the 2D frustrated antiferromagnet. The GGM (vertical) is plotted against the system parameter $\alpha$ (horizontal). Both the quantities are dimensionless.

### C. 2D Frustrated $J_1 - J_2$ Model

We now consider an arrangement of quantum spin$-\frac{1}{2}$ particles on a 2D square lattice, where nearest neighbor spins (both vertical and horizontal) on the lattice are coupled by Heisenberg interactions, with coupling strengths $J_1$, and where all diagonal spins are coupled by the same interactions, with coupling strengths $J_2$. Both $J_1$ and $J_2$ are considered to be positive. The model has attracted a lot of attention [41] due to its connection with high $T_c$-superconductors and its similarity with magnetic materials like $Li_2VOSiO_4$ and $Li_2VOGeO_4$ [42]. Although the different phases of the ground state of this model have been predicted by different numerical as well as approximate analytical methods [43], some debates remain. The Hamiltonian of the system is given by

$$H_{2D} = J_1 \sum_{\langle NN \rangle} \vec{\sigma}_i \cdot \vec{\sigma}_j + J_2 \sum_{\langle \text{diagonals} \rangle} \vec{\sigma}_i \cdot \vec{\sigma}_j, \qquad (9)$$

where $J_1$ and $J_2$ are antiferromagnetic.

In the classical limit, only a first-order phase transition from Néel to collinear at $\alpha \equiv J_2/J_1 = 0.5$ is exhibited by this model. The nature of the phase diagram changes when quantum fluctuations are present, and in this case, the exact phase boundaries are not known. Based on exact diagonalization, series expansion methods, field-theory methods [43], etc., one expects that there are two long range ordered (LRO) ground state phases in the system, that are separated by quantum paramagnetic phases without LRO. These investigations predict that the first transition from Néel to dimer occurs at $\alpha = \alpha_1^c \in (0.35, 0.45)$ while dimer to collinear transition happens at $\alpha = \alpha_2^c \in (0.62, 0.68)$. There are recent experimental observations and proposals of detecting these phases in the laboratory, and they demand a precise quantification of the low temperature phase diagram of this model.

We investigate the behavior of genuine multipartite entanglement of the ground state and study its effectiveness to detect the transitions present in the system. To obtain the ground states, we use exact diagonalization technique as mentioned in the preceding subsection. In Fig 3, we plot the GGM as a function of the driving parameter $\alpha$. From the behavior of the GGM, we predict that the dimer to collinear transition point is in the range $\alpha \in (0.61, 0.62)$ for $N = 16$, and $\alpha \approx 0.81$ for $N = 24$. The indicator here is a maximum appearing between a long-span increase and a long-span decrease in the GGM curve with respect to $\alpha$. Note that there are other short-span increases and decreases (with local maxima in between), which are attributable to finite size errors. The Néel to dimer transition is estimated to occur at $\alpha \approx 0.3$ for both the cases, indicated by a sharp bend there of the GGM curve. Just like in the 1D case, the GGM curves asymptotically go to zero for high $\alpha$. This is because the spin lattice decouples into two spin lattices with nearest neighbor interaction couplings, $J_2$, for very high values of the driving parameter $\alpha$.

### D. The Shastry-Sutherland Model



FIG. 4. The Shastry-Sutherland lattice. The solid lines represent the nearest neighbor interactions with coupling strength $J_1$ between the lattice sites and the ones joined by the dashed lines represent next neighbor interactions with coupling strength $J_2$.

In this section, we study the entanglement properties of systems where the interaction between particles can be modeled by the Shastry-Sutherland Hamiltonian [26]. In the insulators like $SrCu_2(BO_3)_2$, the low-energy spin excitations reside on the spin-half copper ions which lie in two dimensional layers decoupled from each other. The antiferromagnetic exchange couplings between the Cu ions is identical to the Shastry-Sutherland Hamiltonian. The lattice with schematic interactions, for this model, is given in Fig. 4 and the Hamiltonian is given by

$$H_{SS} = J_1 \sum_{NN} \vec{\sigma}_i . \vec{\sigma}_j + J_2 \sum_{NNN} \vec{\sigma}_i . \vec{\sigma}_j. \qquad (10)$$

Here, $J_1 (> 0)$ corresponds to nearest neighbor interaction (indicated by solid lines in Fig. 4) and $J_2 (> 0)$ corresponds to specific next nearest neighbors (indicated by broken lines in Fig. 4). It is known that a simple product of singlet pairs, on the diagonal links, is the ground state of $H_{SS}$ for sufficiently large $\alpha \equiv J_2/J_1$. It has been previously reported [44, 45] that the system undergoes two quantum phase transitions driven by the quantum fluctuations: one is from Néel to an intermediate phase and the other one is from that intermediate phase to dimer. The nature of the intermediate phase is not yet clearly understood [46]. In Fig. 5, we plot the GGM as a function of $\alpha$ for 16 particles on a square lattice interacting via the Shastry-Sutherland Hamiltonian. There are clear signatures of these phase transitions in the figure at $\alpha \approx 1.05$ and at $\alpha \approx 1.53$, as have been predicted by other methods [44, 45]. Due to the computational constraints we are able to report our finding only for $N = 16$. More specifically, we expect that studying the Shastry-Sutherland model requires an exact square structure with even numbers of spins on each side, and hence the next relevant lattice size is $N = 36$. For $N = 16$, we have again used TITPACK ver.2 [40].



FIG. 5. (Color online.) GGM for the Shastry-Sutherland antiferromagnet. GGM (on vertical axis) is plotted with respect to the system parameter $\alpha$ (on horizontal axis) for 16 spins. Both the quantities are dimensionless.

## IV. CONCLUSIONS

An important classification scheme for multipartite entangled quantum states is according to their separability in different partitions. The complexity of such a classification makes it difficult to obtain a unique multiparty entanglement measure, even for pure quantum states. A comparison with the situation for mixed bipartite states is relevant here. While the entanglement of pure bipartite states can be uniquely characterized by a single entanglement measure, a variety of different measures exist for mixed bipartite states. In the case of multiparty quantum states, one can analogously have “pure” and “non-pure” forms of entanglement, corresponding respectively to genuine multiparty entanglement and ones that are not genuine. In this sense, the generalized geometric measure, which is a measure of genuine multiparty entanglement, quantifies the pure form of multiparty entanglement that is present in a multiparty quantum state.

We employed this measure to predict phase diagrams in quantum many-body systems. We began by using the measure to detect the quantum fluctuation driven phase transition in an exactly solvable model, viz. the quantum XY model. We subsequently applied the generalized geometric measure to prototype frustrated quantum spin models, in the quasi two dimensional antiferromagnetic $J_1 - J_2$ model, the two dimensional antiferromagnetic $J_1 - J_2$ model and the Shastry-Sutherland model. The ground states and the corresponding phase diagrams, for the frustrated models, are not known exactly, although there have been several predictions by different methods. We use exact diagonalization techniques to investigate the multipartite entanglement of the ground states of the frustrated models. We find that the genuine multipartite entanglement measure is capable of distinguishing between the gapless and gapped phases in these models.

## ACKNOWLEDGMENTS

R.P. acknowledges support from the Department of Science and Technology, Government of India, in the form of an INSPIRE faculty scheme at the Harish-Chandra Research Institute (HRI), India. We acknowledge computations performed at the cluster computing facility in HRI.

[1] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, Rev. Mod. Phys. **81**, 865 (2009).
[2] S. Sachdev, *Quantum Phase Transistions* (Cambridge University Press, Cambridge, 2011).
[3] M. Lewenstein, A. Sanpera, V. Ahufinger, B. Damski, A. Sen(De), and U. Sen, Adv. Phys. **56**, 243 (2007).
[4] L. Amico, R. Fazio, A. Osterloh, and V. Vedral, Rev. Mod. Phys. **80**, 517 (2008).
[5] S. R. White, Phys. Rev. Lett. **69**, 2863 (1992); S. R. White, Phys. Rev. B **48**, 10345 (1993); G. Vidal, Phys. Rev. Lett. **91**, 147902 (2003); **93**, 040502 (2004); F. Verstraete, D. Porras, and J. I. Cirac, Phys. Rev. Lett. **93**, 227205 (2004); K. A. Hallberg, Adv. Phys. **55**, 477 (2006).
[6] A. Klümper, A. Schadschneider, J. Zittartz, J. Phys. A **24**, L955 (1991); Z. Phys. B **87**, 281 (1992); G. Vidal, Phys. Rev. Lett. **91**, 147902 (2003); D. Perez-Garcia, F.

Verstraete, M.M. Wolf, J.I. Cirac, Quantum Inf. Comput. **7**, 401 (2007).

[7] F. Verstraete and J. I. Cirac, Phys. Rev. A **70**, 060302(R) (2004).

[8] H. Niggemann, A. Klümper, and J. Zittartz, Z. Phys. B **104**, 103 (1997); F. Verstraete and J.I. Cirac, arXiv:cond-mat/0407066; M. Levin and C.P. Nave, Phys. Rev. Lett. **99**, 120601 (2007); H.C. Jiang, Z.Y. Weng, and T. Xiang, Phys. Rev. Lett. **101**, 090603 (2008); Z.C. Gu, M. Levin, and X.G. Wen, Phys. Rev. B **78**, 205116 (2008).

[9] S. Trotzky, Y.-A. Chen, U. Schnorrberger, P. Cheinet, and I. Bloch, Phys. Rev. Lett. **105**, 265303 (2010); M. Cramer, M.B. Plenio, and H. Wunderlich, *ibid.* **106**, 020401 (2011); P.J. Pemberton-Ross and A. Kay, *ibid.*, 020503 (2011), and references therein.

[10] T. Zibold, E. Nicklas, C. Gross, and M.K. Oberthaler, Phys. Rev. Lett. **105**, 204101 (2010); C. Gabriel, A. Aiello, W. Zhong, T.G. Euser, N.Y. Joly, P. Banzer, M. Förtsch, D. Elser, U.L. Andersen, Ch. Marquardt, P.St.J. Russell, and G. Leuchs ibid. **106**, 060502 (2011), and references therein.

[11] M. -F. Yang, Phys. Rev. A. **71**, 030302 (2005)

[12] X.-F. Qian, T. Shi, Y. Li, Z. Song, and C.-P. Sun, Phys. Rev. A **72**, 012333 (2005).

[13] M. N. Bera, R. Prabhu, A. Sen(De), and U. Sen, arXiv:1209.1523 [quant-ph].

[14] A. Shimony, Ann. N.Y. Acad. Sci. **755**, 675 (1995); H. Barnum and N. Linden, J. Phys. A **34**, 6787 (2001); T.-C. Wei and P.M. Goldbart, Phys. Rev. A **68**, 042307 (2003).

[15] M. Balsone, F. Dell'Anno, S. De Siena and F. Illuminatti, Phys. Rev. A **77**, 062304 (2008).

[16] D.A. Meyer and N.R. Wallach, J. Math. Phys. **43**, 4273 (2002).

[17] T.-C. Wei and P. M. Goldbart, Phys. Rev. A **68**, 042307 (2003); T.-C. Wei, D. Das, S. Mukhopadyay, S. Vishveshwara, and P. M. Goldbart, Phys. Rev. A **71**, 060305(R) (2005); T. R. de Oliveira, G. Rigolin, M. C. de Oliveira, and E. Miranda, Phys. Rev. Lett. **97**, 170401 (2006); D. Buhr, M. E. Carrington, T. Fugleberg, R. Kobes, G. Kunstatter, D. McGillis, C. Pugh, D. Ryckman, J. Phys. A: Math. Theor. **44**, 365305 (2011).

[18] A. Sen(De) and U. Sen, Phys. Rev. A **81**, 012308 (2010); A. Sen(De) and U. Sen, arXiv:1002.1253 [quant-ph].

[19] R. Prabhu, S. Pradhan, A. Sen(De), and U. Sen, Phys. Rev. A **84**, 042334 (2011); H.S. Dhar and A. Sen(De), J. Phys. A: Math. Theor. **44**, 465302 (2011); R. Prabhu, A.K. Pati, A. Sen(De), and U. Sen, arXiv:1109.4318 [quant-ph]; H.S. Dhar, A. Sen(De), and U. Sen, arXiv:1110.3646 [quant-ph]; M.N. Bera, R. Prabhu, A. Sen(De), and U. Sen, Phys. Rev. A **86**, 012319 (2012).

[20] G. Misguich and C. Lhuillier, in *Frustrated Spin Systems*, ed. H.T. Diep (World Scientific, Singapore, 2003); C. Lhuillier, arXiv:cond-mat/0502464; F. Alet, A.M. Walczak, and M.P.A. Fisher, Physica (Amsterdam) **369A**, 122 (2006).

[21] M. Rasolt and Z. Tesanović, Rev. Mod. Phys. **64**, 709 (1992); M. Sigrist and T.M. Rice, *ibid.* **67**, 503 (1995).

[22] E. Lieb, T. Schultz, and D. Mattis, Ann. Phys. **16**, 407 (1961); E. Barouch, B.M. McCoy, and M. Dresden, Phys. Rev. **2**, 1075 (1970); E. Barouch and B.M. McCoy, Phys. Rev. **3**, 786 (1971); B.K. Chakrabarti, A. Dutta, and P. Sen, *Quantum Ising Phases and Transitions in Transverse Ising Models* (Springer, Heidelberg (1996)).

[23] C.K. Majumdar and D.K. Ghosh, J. Math. Phys. **10**, 1388 (1969); C.K. Majumdar and D.K. Ghosh, *ibid.* **10**, 1399 (1969).

[24] S.R. White and I. Affleck, Phys. Rev. B **54**, 9862 (1996).

[25] H.J. Mikeska and A.K. Kolezhuk, in *Quantum Magnetism*, ed. U. Schollwöck, J. Richter, D.J.J. Farnell, and R.F. Bishop, (Springer, Berlin, 2004).

[26] B.S. Shastry and B. Sutherland, Physica B+C (Amsterdam) **108**, 1069 (1981).

[27] P.W. Anderson, Science **235**, 1196 (1987).

[28] K. Kim, M.-S. Chang, S. Korenblit, R. Islam, E. E. Edwards, J. K. Freericks, G.-D. Lin, L.-M. Duan, and C. Monroe, Nature (London) **465**, 590 (2010).

[29] J. Struck, C. Olschlager, R. Le Targat, P. Soltan-Panahi, A. Eckardt, M. Lewenstein, P. Windpassinger, and K. Sengstock, Science **333**, 996 (2011); J. Simon, W. S. Bakr, R. Ma, M. E. Tai, P. M. Preiss, and M. Greiner, Nature (London) **472**, 307 (2011).

[30] J. Zhang, M.-H. Yung, R. Laflamme, A. Aspuru-Guzik, and J. Baugh, Nature Communications **3**, 880 (2012).

[31] I. Bloch, J. Dalibard, and W. Zwerger, Rev. Mod. Phys. **80**, 885 (2008); C. Nayak, S.H. Simon, A. Stern, M. Freedman, and S. Das Sarma, *ibid.* **80**, 1083 (2008).

[32] D. Leibfried, R. Blatt, C. Monroe, and D. Wineland, Rev. Mod. Phys. **75**, 281 (2003); H. Häffner, C. Roos, and R. Blatt, Phys. Rep. **469**, 155 (2008).

[33] X.-S. Ma, B. Dakic, W. Naylor, A. Zeilinger, and P. Walther, Nat. Phys. **7**, 399 (2011).

[34] X. Peng, J. Zhang, J. Du, and D. Suter, Phys. Rev. Lett. **103**, 140501 (2009); J. Zhang, T.-C. Wei, and R. Laflamme, *ibid.* **107**, 010501 (2011).

[35] M.A. Nielsen, Phys. Rev. Lett. **83**, 436 (1999); G. Vidal, *ibid.* **83**, 1046 (1999); G. Vidal, J. Mod. Opt. **47**, 355 (2000).

[36] A. Osterloh, L. Amico, G. Falci, and R. Fazio, Nature **416**, 618 (2002); T.J. Osborne and M.A. Nielsen, Phys. Rev. A **66**, 032110 (2002).

[37] M. Matsuda and K. Katsumata, J. Magn. Magn. Mater. **140-145**, 1671 (1995).

[38] R.B. Griffiths, Phys. Rev. **133**, A768 (1964); C.N. Yang and C.P. Yang, *ibid.* **150**, 327 (1966).

[39] F.D.M. Haldane, Phys. Rev. B **25**, 4925 (1982); K. Okamoto and K. Nomura, Phys. Lett. A **169**, 433 (1992).

[40] H. Nishimori, AIP Conf. Proc. **248**, 269 (1991).

[41] P. Chandra and B. Doucot, Phys. Rev. B **38**, 9335 (1988); E. Dagotto and A. Moreo, Phys. Rev. Lett. **63**, 2148 (1989); F. Figueirido, A. Karlhede, S. Kivelson, S. Sondhi, M. Rocek, and D. S. Rokhsar, Phys. Rev. B **41**, 4619 (1990); N. Read and S. Sachdev, Phys. Rev. Lett. **66**, 1773 (1991); H.J. Schulz and T. A. L. Ziman, Europhys. Lett. **18**, 355 (1992); R. Darradi, O. Derzhko, R. Zinke, J. Schulenburg, S.E. Krüger, and J. Richter, Phys. Rev. B **78**, 214415 (2008), and references therein.

[42] R. Melzi, P. Carretta, A. Lascialfari, M. Mambrini, M. Troyer, P. Millet, and F. Mila, Phys. Rev. Lett. **85**, 1318 (2000); H. Rosner, R. R. P. Singh, W. H. Zheng, J. Oitmaa, S. L. Drechsler, and W. E. Pickett, Phys. Rev. Lett. **88**, 186405 (2002); R. Nath, A.A. Tsirlin, H. Rosner, and C. Geibel, Phys. Rev. B **78**, 064422 (2008); T. Yildirim, Phys. Rev. Lett. **101**, 057010 (2008); Q. Si and E. Abrahams, Phys. Rev. Lett. **101**, 076401 (2008).

[43] J. Richter and J. Schulenberg, Eur. Phys. J. B **73**, 117 (2010); J.K. Kim and M. Troyer, Phys. Rev. Lett. **80**, 2705 (1998); T. Pardini and R.R.P. Singh, Phys. Rev. B

**79**, 094413 (2009).

[44] C. H. Chung, J. B. Marston and S. Sachdev, Phys. Rev. B. **64**, 134407 (2001).

[45] M. Albrecht and F. Mila, Europhys. Lett. **34**, 145 (1996).

[46] W. Zheng, J. Oitmaa, and C. J. Hamer, Phys. Rev. B. **65**, 014408 (2001); M. A. Hajj and J. Malrieu, Phys. Rev. B. **72**, 094436 (2005); R. Darradi, J. Richter, and D. J. J. Farnell, Phys. Rev. B. **72**, 104425 (2005); R. W. Chhajlany, P. Tomczak, A. Wójcik, and J. Richter, Phys. Rev. A. **75**, 032340 (2007); S. Moukouri, Phys. Rev. B. **78**, 132405 (2008).